

IS IT POSSIBLE THAT FOOD RICH IN CALORIES PREDISPOSES TO HEART ATTACK?

Chibisov S.¹, Singh R.², Eremina I.¹, Vargova V.³, Pella D.³,
Otsuka K.⁴, Savrova O.¹, Kharlitzkaya E.¹ and Agarval R.⁵

*1 Faculty of Medicine, The People's Friendship University of Russia,
Moscow, Russia*

2 The Halberg Hospital and Research Institute, Moradabad, India

3 Faculty of Medicine, The Safaric University, Kosice, Slovakia

4 The Tokyo Women's Medical University Medical Centre, Tokyo, Japan

5 Network of Young Doctors and Health Administrators, Moscow, Russia

Acute myocardial infarction (AMI) is a highly dynamic event, which is associated with marked neuroendocrinological dysfunction apart from heart damage. The immediate trigger for heart attack is not known exactly. There is a marked increase in sympathetic activity, oxidative stress and deficiency of minerals as well as antioxidants, during heart attack. Clinical studies have reported an increased incidence of reinfarction, sudden death, coronary constriction, myocardial ischaemia and angina, during first quarter of the day when there is rapid withdrawal of vagal activity and increase in sympathetic tone.

Keywords: large meals, breakfast, trans fatty acids, cytokines, lipoproteins, glucose.

Introduction

In one case control study, among 202 patients of heart attack, there was a significant ($P < 0.02$) increase in cardiac events in the second quarter of the day (6-12 hours) compared to other quarters respectively (16.8%, 41.0%, 13.8%, 28.2%). This characteristic remained common in both men and women and among patients with and without known heart attack ($n=52$), diabetes ($n=53$) and hypertension ($n=75$). Triggers of heart attack were noted among 162 (82.2%) of the patients. Brain related and psychological mechanisms were; emotional stress (45.5%), sleep deprivation (27.7%), cold climate (29.2%), hot climate (24.7%), large meals (47.5%) and physical exertion (31.2%). These triggering factors are known to enhance sympathetic activity and decrease vagal tone, resulting into increased secretion of plasma cortisol, noradrenaline, aldosterone, angiotension converting enzyme, interleukin-1,2,6,18 and tumor necrosis factor-alpha, that are pro-inflammatory. There is also a deficiency in the serum levels of vitamin A, E, C, coenzyme Q10 and magnesium, potassium, melatonin, interleukin-10 (anti-inflammatory) and increase in TBARS,

MDA and diene conjugates, TNF-alpha and IL-6 which are indicators of oxidative damage and pro-inflammatory respectively. It is not clear whether the predisposition of ACS is due to size of the meals or proinflammatory content of meals.

The exact mechanism and the immediate triggers for acute heart attack and brain attacks are not known. Clinical manifestations of these attacks also do not occur at random times but according to a time structure [1-7]. It is possible that certain external activities, known as triggers, play a major role in the occurrence of these attacks. In one study [5], half of all heart attack patients reported a temporal relationship between characteristic activities and occurrence of the attack. Emotional stress, sleep deprivation, large meals, mild and heavy physical exertion were the most frequently reported triggering factors of heart attacks in various studies [1-7, 11-13].

There appears to be a powerful evidence of a link between these triggers and the sequences of cellular and pathophysiological events that are postulated as being responsible for poor supply of blood in the heart. The morning increase in platelet aggregability is a

most frequent trigger of heart attack, which may be the result of increased secretion of cortisol, aldosterone, catecholamines, angiotensin, free fatty acids, triglycerides [7-11] and decreased vagal tone that are known to be enhanced by food and benefited by fasting. Apart from above triggers, presence of risk factors of low heart rate variability, such as excess fat, more insulin resistant, obesity and pollution may also act by enhancing triggers and trigger-induced brain-related and hormonal mechanisms in the development of poor supply of blood to heart [1]. Both clinical and biochemical factors may be related to a molecular clock present in the brains suprachiasmatic nucleus.

The triggers of acute coronary syndromes

Heart attack is a highly dynamic event, which is associated with marked neuroendocrinological dysfunction apart from heart damage. The immediate trigger for heart attack is not known exactly. Studies conducted by various investigators [2-7, 11-13], have demonstrated, a marked increase in sympathetic activity, oxidative stress and a decrease in parasympathetic activity in association with increase in catecholamines, cortisol and decrease in 1). melatonin [1]. In one case control study by Singh et al [5], among 202 patients of ACS, brain related and psychological mechanisms were observed as given : emotional stress (45.5%), sleep deprivation (27.7%), cold climate (29.2%), hot climate (24.7%), large meals (47.5%) and physical exertion (31.2%). These triggering factors are known to enhance sympathetic activity and decrease vagal tone, resulting into adverse biochemical environment in the body tissues. In another study, by Singh et al, among 54 patients, that were included in this study, 41 patients had full heart attack, 5 patients possible 2nd stage, 8 cases first stage. The control subjects (n=85) were randomly selected from the population of the city of Moradabad drawn from a similar age range of subjects after exclusion of heart attack (n=9), diabetes(n=6) and excess intake

of trans fatty acids (n=20). Large breakfast was a predisposing factor of heart events in the second quarter of the day and it was significantly associated with metabolic reactions. The findings indicate that acute reactions as a result or as circadian rhythms appear to be important in the pathogenesis of complications in heart attack and that a large breakfast in association with nitrite deficiency, may further trigger the circadian rhythms. However more studies in a large number of subjects would be necessary to confirm our findings. The TRIMM [6] and the MILIS study [3] also showed that large meals is a trigger for ACS.

Circadian rhythmicity

Clinical studies have reported an increased incidence of attacks, sudden death and poor blood supply to heart, during first quarter of the day when there is rapid withdrawal of vagal activity and increase in sympathetic tone. In one case control study by Singh et al [5], among 202 patients of heart attack, there was a significant ($P < 0.02$) increase in cardiac events in the second quarter of the day (6-12 hours) compared to other quarters respectively (16.8%, 41.0%, 13.8%, 28.2%). This characteristic remained common in both men and women and among patients with and without known heart attack (n=52), diabetes (n=53) and hypertension (n=75). In another study, by Singh et al [4] among 54 patients with ACS, there was a significant greater incidence of cardiovascular events in the second quarter of the day (6.00 AM to 12.00 noon) compared to 3rd quarter. In another study, by Singh et al [4] among 54 patients with ACS. Circadian rhythmicity of cardiovascular events have also been reported in the TRIMM study as well as in the MILIS Study.

Biochemical mechanisms

There is evidence that there is increased secretion of plasma cortisol, noradrenaline, aldosterone, angiotensin converting enzyme, interleukin-1,2,6,18 and tumor necrosis factor-alpha, that are pro-

inflammatory during ACS [1-7]. There is also a deficiency in the serum levels of vitamin A, E, C, coenzyme Q and magnesium, potassium, melatonin, interleukin-10 (anti-inflammatory) [4, 5, 12, 13]. In our studies [4, 5, 7], we found a decrease in magnesium, potassium, vitamin A, E, C and beta carotene and increase in TBARS, MDA and diene conjugates, TNF-alpha and IL-6 which are indicators of oxidative damage and pro-inflammatory, respectively. Mean lipoprotein(a), total cholesterol and triglycerides were significantly higher and mean nitrite level, lower in the ACS group, compared to control group [5]. Lp(a), triglycerides, blood glucose, plasma insulin, malondialdehyde, diene conjugates, TBARS and TNF-alpha and IL-6 which were significantly greater during acute phase, showed a significant decline, and serum nitrite and coenzyme Q, an increase, at 4 weeks of follow up, when the acute reactions evoked by ACS, were controlled [4].

Large meals and acute coronary syndromes

There is little information on the role of nutrition in the onset of acute reactions and ACS. Several workers observed that subjects consuming large breakfast and meals especially dinner were more common victims

of attacks, compared to apparently healthy subjects of same age and sex [2-6]. Higher intake of hydrogenated fat was also more common among ACS patients in one study which is known to have adverse effects of lipids and inflammation(4). It is known that different types of fatty acids and refined carbohydrates in the diet, influence different physiologically relevant mechanisms, especially those concerned with haemostasis and inflammation in the body [19-21].

As the role of various factors, influencing clot formation in the coronary artery and thrombosis on risk of heart and stroke are more firmly established, our knowledge concerning the effects of different types of nutrients on these factors remains limited. A recent study, showed that postprandial leptin response was lower after a carbohydrate meal in obese women, than in lean controls, suggesting an impairment of such response of leptin regulation due to obesity. It is known that postprandial leptin changes are higher after carbohydrate meal than after a fat meal similar to insulin changes.

We found that consumption of a large breakfast was associated with greater pro-inflammatory cytokines; TNF-alpha and IL-6, than in subjects consuming low energy breakfast (Table 1, 2).

Table 1. Food intakes among patients and control subjects (reference 4)

| Foods intake (g/day) | Before acute coronary syndrome (n=54) | Controls (n=85) |
|------------------------------|---------------------------------------|-----------------|
| Antiatherogenic foods(g/day) | | |
| Wheat, rice and millets | 345 ± 43.3* | 486 ± 69.6 |
| Roots and tubers | 49.5 ± 4.2 | 52.1 ± 5.2 |
| Fruits and vegetables | 161 ± 13.2* | 221 ± 25.4 |
| Legumes and pulses | 35 ± 6.7* | 53 ± 9.1 |
| Almonds and walnuts | 0.5 ± 0.2* | 1.5 ± 1.2 |
| Fish | 10 ± 5.9* | 16.7 ± 6.6 |
| Mustered oil | 10 ± 4.4* | 18 ± 6.9 |

| Foods intake (g/day) | Before acute coronary syndrome (n=54) | Controls (n=85) |
|---|---------------------------------------|-----------------|
| Proatherogenic foods(g/day) | | |
| Butter, Indian ghee, hydrogenated oils. | 38.0 ± 10.0** | 16.7 ±5.9 |
| Oils rich in omega-6 fatty acids | 22.5 ± 4.7* | 11.8 ± 3.5 |
| Total visible fat | 70.5 ± 14.6* | 46.5 ± 11.7 |
| Sugar, bread, biscuits | 208± 57.8** | 115± 29.6 |
| Meats(chicken, beef, gote, eggs) | 27± 10.7** | 40± 12.8 |
| Milk and it products | 152 ± 34.6* | 112 ± 22.5 |

Values are mean(standard deviation). *=P <0.05, ** P<0.01; P values were obtained by Student, s t test for continuous variables.

Table 2. Association of size of meals with biochemical risk factors of coronary syndromes.

| Biochemical data | Large breakfast, n=22 | | Small breakfast (n=32) | |
|-------------------------|-----------------------|---------------|------------------------|---------------|
| | Baseline | After 4 weeks | Baseline | After 4 weeks |
| Triglycerides (mmol/L) | 1.88±0.61 | 1.70±0.38* | 1.81+0.60 | 1.64+0.32* |
| Blood glucose (mmol/l) | 7.7±1.6 | 6.0±1.2* | 6.6+1.4* | 5.5+0.30* |
| Plasma insulin(mg/dl) | 47.5+11.3 | 36.3+5.6** | 43.2+8.8* | 27.6+3.5* |
| TBARS(pmol/l) | 1.87+0.46 | 1.32+0.33* | 1.77+0.42 | 1.30+0.31* |
| Malondialdehyde(pmol/l) | 2.68+0.34 | 2.02+0.21* | 2.66+0.33 | 2.01+0.21* |
| Diene conjugate(OD) | 27.5+4.2 | 24.6+4.0* | 26.2+4.1 | 24.2+3.5* |
| Coenzyme Q10(ug/ml) | 0.21+0.02 | 0.32+0.23* | 0.23+0.03 | 0.45+0.24* |
| Interleukin-6(pg/ml) | 32.6+6.2 | 22.5+4.3* | 27.5+5.2* | 20.6+0.22* |
| TNF-alpha(ug/dl) | 42.5+12.8 | 23.6+4.1* | 38.2+10.6* | 19.6+0.18* |
| Lipoprotein(a) Ug/ml | 23.1±5.4 | 20.1±4.2* | 22.5+4.6 | 19.7+4.1* |

Values are mean standard deviation.*= P<0.05,**=P< 0.01.P values were obtained by analysis of variance by comparison of large breakfast verses small breakfast groups.

Increased levels of TNF-alpha and IL-6 have been observed in patients with obesity, type 2 diabetes insulin resistance, glucose intolerance, hypertriglyceridemia, hyperleptinemia and heart attack, which could be greater in patients with attacks. There is no previous evidence that people consuming large breakfast have greater proinflammatory cytokines, as observed in our study. There is potential evidence, however, that ACS may be associated with greater levels of proinflammatory

cytokines due to metabolic reactions such as catecholamines, cortisol, serotonin, hyperinsulinemia, hypertriglyceridemia and hyperglycemia that are common in patients with attacks. It is possible that people consuming large, fatty breakfast enhance the release of catecholamines, glucose, insulin, triglycerides which may have an adverse effects on cardiovascular function.

However, A few studies just published [14-18] provides added evidence that feeding

at the “wrong time” can lead to weight gain. These recent findings on mice were extended to humans in 1973 [17]. A start with nutraceuticals indicates that in a person investigated by varying the administration times of coenzyme Q10 (CoQ10), the circadian stage played an important role [14, 18]. It is possible that a heavy breakfast as well as a heavy dinner, both may have adverse effects.

Experiments in animals indicate that ventromedial hypothalamic (VMH) lesion in the brain, in rats induces hyperphagia, and excessive weight gain, fasting hyperglycemia, hyperinsulinemia, hypertriglyceridemia and glucose intolerance [19]. There may be suppressed splenic natural killer cell activity, when the animals are hyperphagic and obese. There is increased release of interferon-alpha, IL-1, IL-2 and TNF in response to non-inflammatory and inflammatory stresses. Longterm infusion of norepinephrine plus serotonin into the VMH part of brain, impaired pancreatic islet cell function in as much as these abnormalities are observed in insulin resistant animals by several researchers [19]. It is possible, that heart attack is associated with high level of psychological and hormonal stress, which may have caused the release of proinflammatory cytokines which can damage heart and endothelial cells resulting into increased susceptibility to clot formation and ACS and heart failure.

Adverse effect of diet

Dietary fat composition can also influence platelet function and platelet aggregation and blood clotting [25]. Recent evidence indicate that dietary butter or coconut, increase the sensitivity of platelets to aggregation and clot formation and enhances the release of harmful catecholamines [25]. Hypercholesterolemia is known to be associated with an increased sensitivity of platelets to aggregating agents. A large breakfast rich in fat may enhance platelet aggregation and clotting resulting into heart attack, as noted in our study. Reduction in the saturated fat in the diet may decrease, platelet aggregation and catecholamines resulting into decrease in

blood pressure and heart rate. A diet rich in n-3 fatty acids may decrease cytokines and hence may be beneficial because it is cardioprotective as well as neuroprotective [19-25].

Role of fasting or low energy indo-mediterranean diet

We have found that subjects eating low energy diet had low blood pressure, lower glucose, insulin, cholesterol, triglycerides and cytokines. In 32 subjects, on no breakfast, fruits and vegetables in the lunch and normal dinner, we observed a very low blood pressure no vascular variability and no overweight and obesity, compared to subjects eating normal meals. It is possible that no breakfast or small amount of breakfast, and small frequent meals including dinner containing protective foods such as almonds, walnuts and black raisins, whole grains bread, legumes, fruits, vegetables and salads, 5-6 times in a day, may have a beneficial effect on the harmful body chemistry that occurs during the 2nd and 4th quarters of the day [25]. It is suggested that take small breakfast and dinner and normal lunch and supper to avoid the adverse effects of foods during circadian rhythms of our body and prevention of heart attacks. Recent studies also indicate that fasting for few hours is also cardio and neuroprotective.

References:

1. Singh RB, Cornelissen G, Weydahl A, Schwartzkopft O, Katinas G, Otsuka K et al, **Circadian heart rate and blood pressure variability considered for research and patient care.** *Int J Cardiol* 2003,87:9-28.
2. Muller JE, Befler GH, Stone PH. **Circadian variations and triggers of onset of acute cardiovascular disease.** *Circulation* 1989,4: 733-40.
3. Muller JE, Stone PH, Turi ZG, Rutherford JD, Czernsler CA, Parker C, et al. **The MILIS Study group. circadian variations in the frequency of onset of acute myocardial infarction.** *N Engl J Med* 1985,313:1315-22.
4. Singh RB, Pella D, Neki NS, Chandel JP, Rastogi S, Mori H et al. **Mechanism of acute myocardial infarction study (MAMIS).** *Biomed Pharmacol*, 2004, 58:(Supple)111-115.

5. Singh RB, Pella D, Sharma JP, Rastogi S, Kartikey K, Goel VK, Sharma R, Neki NS, Kumar A, Otsuka K. **Increased concentrations of lipoprotein (a), circadian rhythms and metabolic reactions, evoked by acute myocardial infarction, associated with acute reactions, in relation to large breakfasts.** *Biomed Pharmacother* 2004, 58: (Supple)116-122.
6. Willich SN, Loewel H, Lewis M et al. (TRIMM Study group). **Association of wake time and the onset of myocardial infarction.** *Z Kardiol* 1991; 80 (Supple 3) 105-112.
7. Singh RB, Niaz MA, Rastogi SS, Sharma JP, Kumar R, Bishnoi I, Begom R. **Plasma levels of antioxidant vitamins and free radical stress in patients with acute myocardial infarction.** *Acta Cardiol*, 1994, 49:411-52.
8. Das UN. **Free radicals, cytokines and nitric oxide in cardiac failure and myocardial infarction.** *Mol Cell Biochem* 2000,215:145-152.
9. Singh RB, Neki NS, Kartikey K, Pella D, Kumar A, Niaz MA et al. **Effect of coenzyme Q10 on risk of atherosclerosis in patients with recent myocardial infarction.** *Mol Cell Biol Chem*, 2003, 246:75-82.
10. Singh RB, Cornelissen G, Siegelova J, Homolka P, Halberg F. **About half weekly pattern of blood pressure and heart rate in men and women of India.** *Scripta Medica* (BRNO), 2002, 75:125-128.
11. Singh RB, Niaz MA, Cornelissen G, Otsuka K, Siegelova J, Fiser B, Halberg F. **Circadian rhythmicity of circulating vitamin concentrations.** *Scripta Medica* (BRNO),2001,74:93-96.
12. Singh RB, Weydahl A, Otsuka K, Watyanabe Y et al. **Can nutrition influence circadian rhythm and heart rate variability.** *Biomed Pharmacol* 2001, 55 (Supple) 115-124.
13. Reis F, Hermida RC, Sonza I, Maldonado J, Tavares P, Fontes-Reibeiro CA, Teixeira HM, Alcobia T, Almeida I, Teixeira F. **Circadian seasonal variation of endogenous ubiquinone plasma level.** *Chronobiol Interna* 2002,19:599-614.
14. Franz Halberg, Germaine Cornelissen and R.B.Singh. **Timing nutraceuticals?** *The Open Nutra J*. (In press)
15. Arble DM, Bass J, Laposky AD, Vitaterna MH, Turek FW. **Circadian timing of food intake contributes to weight gain.** *Obesity* 2009; doi:10.1038/oby.2009.264 (advance online publication 3 September 2009).
16. Arble DM, Goldschmidt C, Vitaterna MH, Turek FW. **Circadian eating and sleeping patterns in the night eating syndrome.** *Sleep* 2009; 32: A61.
17. Halberg F. **Protection by timing treatment according to bodily rhythms: an analogy to protection by scrubbing before surgery.** *Chronobiologia* 1974; 1 (Suppl. 1): 27-68.
18. Halberg F, Haus E, Cornelissen G. **From biologic rhythms to chronomes relevant for nutrition.** In: **Marriott BM, editor. Not Eating Enough: Overcoming Underconsumption of Military Operational Rations.** *Washington DC: National Academy Press;* 1995. p. 361-372. <http://books.nap.edu/books/0309053412/html/361.html#pagetop>
19. Dube MG, Xu B, Kalra PS, Sninsky CA, Kalra SP. **Disruption in neuropeptide Y and leptin signaling in obese ventromedial hypothalamic – lesioned rats.** *Brain Res* 1999, 816:38-41.
20. Dube MG, Kalra PS, Crowley WR, Kalra SP. **Evidence of a physiological role for neuropeptide Y in ventromedial hypothalamic lesion-induced hyperphagia.** *Brain Res* 1995, 690:275-81.
21. Takahashi A, Ishimaru H, Ikarashi Y, Maruyama Y. **Aspects of hypothalamic neuronal systems in VMH lesion induced obese rats.** *J Auto Nerv Syst* 1994, 48:213-18.
22. Ohtani N, Ohta M, Sugano T. **Microdialysis study of modification of hypothalamic neurotransmitters in streptozotocin-diabetic rats.** *J Neurochem* 1997, 69:1622-26.
23. Funahashi T, Shimomura I, Hiraoka H, et al. **Enhanced expression of rat obese(ob) gene in adipose tissues of ventromedial hypothalamus (VMH) lesioned rats.** *Biochem Biophys Res Commun* 1995, 211:469-73.
24. Singh RB, Pella D, Otsuka K. **Can brain dysfunction be a predisposing factor for metabolic syndrome?** *Biomed Pharmacother J*, 2004, 58:(Supple)58-68.
25. Singh RB, DeMeester F, Mechirova V, Pella D, Otsuka K. **Fatty acids in the causation and therapy of metabolic syndrome.** In **Wild Type Foods in Health Promotion and Disease Prevention**, editors Fabien De Meester and RR Watson, *Humana Press, NJ* 2008, p 263-284.