

## GEOLOGICAL AND GEOCHEMICAL PROPERTIES OF OIL SHALE IN AZERBAIJAN AND PETROLEUM POTENTIAL OF DEEP-SEATED EOCENE-MIOCENE DEPOSITS

Abbasov O.R.

*Institute of Geology and Geophysics of Azerbaijan National Academy of Sciences,  
Baku, e-mail: ortal80@bk.ru*

The paper is devoted to a detailed study of oil shale that is considered a new alternative energy and fuel resource for Azerbaijan. More than 60 surface manifestations of oil shale, related to sediments between Eocene and Miocene in the territories Shamakhi-Gobustan, Absheron, Pre-Caspian-Guba, Vandam-Lahij and etc. of the Republic are investigated on its distribution regularity and geochemical properties. Oil shale of these epochs, which can't be reached due to exploration wells, but brought from the different depths to the Earth's surface (where oil shale surface manifestations widely distributed) by eruptions of mud volcanoes have been studied as well. Along with geochemical study of oil saturated rocks, found in ejected production of mud volcanoes, a comparative analysis have been conducted on heavy fractions, obtained from these rocks and oil shale. The results of these studies show that bitumen (its heavy fractions) of Eocene-Miocene oil shale and oil-saturated rocks are similar, and may cause to formation and accumulation of hydrocarbons in ultra-deep sediments of mud volcanic areas. All conducted investigations are increasing the topicality of perspectives of shale gas in Azerbaijan, which is non-traditional for the country.

**Keywords:** Azerbaijan, distribution regularity, geochemistry, organic matter, oil shale, shale gas

The rich hydrocarbon potential of Azerbaijan has a great influence in the status of country. Integrated studies are expanded for further enrichment of hydrocarbons reserves of Republic. To explore new oil and gas fields in deeper sediments of onshore and offshore territories becomes very relevant. However, the energy potential of the country should not depend only on oil and gas resources, and together with them the Republic is rich in non-traditional alternative sources of energy, such as natural bitumen, oil shale and gas hydrates [1, 11].

Currently, a number of countries are widely used of oil shale to produce shale gas (US, Canada, China and etc.), cement (Estonia, China, Germany), agricultural fertilizers (Estonia, Switzerland), burn at power plants (Estonia, China, Germany, etc.), obtain medical (France, Russia, China) and chemical products (China, Estonia, Russia) and etc.

Researches on oil shale of Azerbaijan have been carried out at the last century [4, 9, 10]. More detailed studies have been carrying out since 2000 by scientists of Institute of Geology and Geophysics of the Azerbaijan National Academy of Sciences. Employees of Department "Mud Volcanism" of the Institute have been studying geological, geochemical properties and probably reserves of oil shale [1, 2, 6, 7, 11, 12].

### Distribution regularities of oil shale in Azerbaijan

The most widespread areas of oil shale in the Republic are observed in Shamakhi-Gobustan, Absheron, Pre-Caspian-Guba and Vandam-Lahij and etc. regions (Fig. 1). There are more than 60 surface manifestations of oil shale in these regions, distributed in a wide stratigraphic range (from the Cretaceous to Miocene).

### Shamakhi-Gobustan region

The region is located in the south-eastern part of the Greater Caucasus and has a very complex geological and tectonic structure. There are about 120 mud volcanoes, more than 30 manifestations of oil shale, as well as some oil and gas fields in the region. The known oil shale manifestations are developed mainly in the Central Gobustan and Shamakhi tectonic zone.

The geological structure of the region consists of Mesozoic-Cenozoic sediments [12]. In generally, related to Goradil-Masazir Fault Zone and Gujur-Gyzyldash Thrust, three blocks: north allochthonous, central para-autochthonous and southern autochthonous are separated in Gobustan (Fig. 2).

Oil shale of the Northern Gobustan belonging to the Cretaceous deposits, have no commercial value [2].

Many of oil shale manifestations are observed in areas of the Central Gobustan associated with Paleogene-Miocene sediments. The Central Gobustan covers areas, relating to para-autochthonous tectonic block (Bayanata), which indicates favorable paleogeographic and paleotectonic conditions of the zone for the formation of oil shale. Paleogene-Miocene sediments are involved in the geological structure of Bayanata block, have thickness of 2,5–4,5 km. In contrast to these structural and facies complexes, deposits of the same age, observed in north of Geradil-Masazir fault zone (north allochthonous) are located under the Cretaceous sediments (consisting of flyshoids). It is assumed that the sediments of Paleogene-Miocene age in Bayanata block have been compressed between the carbonate complexes (Upper Cretaceous age) of allochthonous and the sediments (Cenozoic age) of autochthonous,




Fig. 1. Schematic map of oil shale distribution regions in Azerbaijan




Fig. 2. Geological map of Gobustan [5]

increasing up to 11 km. In such a complex tectonic environment, there were favorable facies and paleogeographic (accumulation of organic matters in silt sediments, existence of shallow, broad bays and continental lagoons) conditions for the accumulation of organic compounds, which forming oil shale. The sediments accumulated with short geochronological breaks and at subsequent stages, a result of dynamic metamorphism these exposed to the process of oil shale formation.

According to the spatio-temporal distribution, the development of oil shale in the Central Gobustan are associated with complex tectonic structure of the region and accumulation of organic compounds with breaks, started from the Middle Eocene, continued in the Maikopian and Konk, ended in the Meotian during the process of lithogenesis.

In general, the number of oil shale manifestations in Gobustan and their probably resources are reduced in the direction from the center to the south-east [1] (Fig. 3).

per Maikopian, Konk and Meotian. The exceptions are related to two manifestations (Goytapa and Uchtepe), the age of Eocene, are located on the border with Shamakhi-Gobustan region.

Conditional border between clay and sandy-clay facies of the Upper Maikopian, traced in the direction of the north wing of Shorbulag folds over of mountains Garaheybat and Bozdag-Qobu, and further to the southeast to Puta Cape. Clay facies of the Upper Maikopian developed in the northern, north-western parts of the Absheron Peninsula and linked to the nature of folding – diapirism occurrence, composing the crest of folds. The most representative section is Riki horizon, situated in the northern slope of Mountain Uchtepe-Shorchala, with thickness of 112 m, composed layers of dark, chocolate-brown clay and black oil shale. The last traced in sections of Mountains Goytepe, Orjandag and Fatmai.

Thus, a broad band of oil shale in the section of Riki horizon coincides with the deepest part of the Upper Maikopian Basin and extends



Fig. 3. Map of probable resources of oil shale in Shamakhi-Gobustan region (scale 1:1,000,000).

Oil shale manifestations:

- 1 – Gizmeydan; 2 – Talishnuru; 3 – Khilmilli; 4 – Angakharan; 5 – Shamakhi;
- 6 – Arabshalbashi; 7 – Jeirli; 8 – Agdara; 9 – Garaja; 10 – Ahudag; 11 – Baygushlu;
- 12 – Shaiblar; 13 – Tuva; 14 – Shahandag; 15 – Kichik Siyaki; 16 – Garayokhush;
- 17 – Garigishlag; 18 – Alagishlag; 19 – Baygushgaya; 20 – Jangichay; 21 – Jangidag;
- 22 – Boyuk Siyaki; 23 – Gibladag; 24 – Kechallar; 25 – Sungur; 26 – Bayanata; 27 – Girdag;
- 28 – Saridag; 29 – Pirekeshkyul; 30 – Islamdag; 31 – Agburun

### Absheron region

The geological structure of Absheron consists of clastic-carbonate rocks of the Upper Cretaceous and Cenozoic. There are more than 15 surface manifestations of oil shale in the region, relate mainly to the sediments of the Up-

per Maikopian, Konk and Meotian. The exceptions are related to two manifestations (Goytapa and Uchtepe), the age of Eocene, are located on the border with Shamakhi-Gobustan region.

Oil shale of Konk horizon developed within the Shorbulag and Garaeybat areas in the Western Absheron, has a thickness of 30 m. Whitish foliated oil shale is traced in



Uchtepe-Ilkhdag syncline and 2 km north-west of station Guzdek along the south-eastern end of Geytepe folds. Here, 120 m thickness of Konk-Karagan sediments are observed as a pack of gray, brown-gray foliated oil shale, with a thickness of 18,7 m. Within the southern pericline of Kechaldag- Zigilpiri fold, has been found Karagan-Konk sediments with frequent interlayers of oil shale, traces for several kilometers: in a section, near the Mountain Zigilpiri with a thickness of 26 m, allocated 7 layers of gray, dark gray oil shale. In addition, they are also recorded to south of the village Masazir and north of the mud volcano Kechaldag, in the western part of the Lake Shirinnour in a distance of 1,5 km. Further to east from station Binagadi, oil shale have been found in Karagan-Konk sediments, have a thickness of 125–130 m [6].

In the north-western and northern parts of Absheron Peninsula, oil shale is observed in sections of the north-eastern wing of Fatmai, Orjandag and Saray folds. Total thickness of Konk sediments in the area is 20–25 m. In Absheron Peninsula, oil shale is traced in section of Meotian, spreads throughout the area and characterizes by different thicknesses.

In the southern part of West Absheron, oil shale is observed in Shorbulaga area, Kosmalidag synclines and Uchtepe, as well as and in a considerable part of the hill Damlamaja down to the town Garaeybat. The others distributions of oil shale are traced in the areas of Ateshgah, Khirdalan, Shabandag, Binagadi and to the north of the mud volcano Keyreki, near Kerpyukshor syncline [10] with 1,5 km length and 4,5 m thickness to the west of station Masazir and district Guzdek. In the northern part of Absheron, oil shale found along the right bank of Sumgaitchay River, in the northern Geytepe Mountain, western and eastern periclinal of Jorat folds, north-east wing of Orjandag (thickness of 22 m, length of 650 m), Saray (thickness of 6 m, length more than 1 km) and Fatmai folds. The thickness of Meotian in these areas ranges from 72 to 130 m.

#### **Pre-Caspian-Guba region**

Related to Upper Cretaceous sediments, oil shale are mostly located in the north-east wing of Zarat syncline. The most studied section is in Atachay River, near the village of Bakhishli in Khizi region. The section consists of dark gray bituminous marl, black oil shale and the total thickness is 27 m. The bituminous oil shale also has been found in the section of the Lower Cretaceous (Albian), in the south-eastern pericline of Kemchi fold with the thickness of 3,5 m in Altigaj area.

The Eocene sediments are presented in two facies in the area: bituminous marl in the north-

west (from Samur River to the city of Shabran) and clay, clay shale in the south-east. The thickness of Middle Eocene in this layer is not more than 40–50 m. Here, there are basically no significant oil shale manifestations with much practical value. Also, minor interlayers of oil shale encountered in some sections of the wells (Siyazan, Saadan and Shuraabad) [4].

The layers of Maikopian series (Oligocene-Upper Miocene) oil shale found in the north-west of Pre-Caspian-Guba region do not attract the attention of the economic prospect.

Oil shale of Konk sediments observes in the Chandagar coast with thickness of 8 to 15 m, but Meotian sediments only in the south-east of Shuraabad district. In the Yashma area, two layers of oil shale are separated in the section of Meotian.

#### **Guba oil shale deposit**

Guba oil shale deposit is located 25 km south of Guba city. Oil shale of deposit relates to Upper Sarmatian, ranges from 27 to 255 m and alternates with layers of clay shale, which hardly differ from oil shale. The greatest practical interest is associated with the segment of the Upper Sarmatian layer, length of 29 km, elongated in the NW-SE direction from Gudiyalchay River to Velvelichay River (Fig. 4), although the individual manifestations of oil shale are found to Gilgilchay River. To the north-west of Gudiyalchay River, oil shale appears like an individual spots in the watershed Gusarchay-Tahirjalchay and village Anig [7]. In this direction, the overall thickness of the upper Sarmatian increases up to 1350 m and transgressive bedding of oil shale also traces in the Middle and Lower Sarmatian. Sheet black oil shale in the section has a monoclinical bedding.

Larger manifestations of oil shale have been studied in the area between rivers Velvelichay and Garachay. There are three areas that contain oil shale with the most favorable properties for practical use. The first area, a length of 4,7 km, includes 11 layers of oil shale. The second area, situated between rivers Kamalchay and Chagachukchay, a length of 3 km, includes 14 layers. The third area, 1,5 km, located on the right bank of Garachay River contains 16 layers of oil shale (Fig. 5).

#### **Diyally oil shale deposit**

The deposit is located 7 km east of Ismailly city, 1,5 km the northeast of Diyally village, in strong cross wooded area, at an altitude of 800 m. Tectonically point view, the area relates to the complicated zone of Vandam Lahij where at the Upper Sarmatian time there were favorable geological and geochemical conditions for the formation of oil shale. The main structural

element is tilted to the south anticlinal fold of the northwest with a torn stretch to fault the southern wing, with dips 50–55°. The core of the folds in the west is composed of Upper Cretaceous rocks, in the south-east of the Maikopian and Upper Sarmatian sediments. North of these folds at a distance of 1,5 km there are two troughs (moulds), relates to Sarmatian sediments, and which belonging to Diyally oil shale deposit. The thickness of oil shale layer varies 300–370 m, in the north is hidden under limestone thrust of Kemchi suite. At the base of the section of the Upper Sarmatian, lies pack of basal conglomerates, with a thickness of up to 70 m, under which a pack of layered clays with interbedded sandstone and oil shale. The last one confined to the upper half of the sec-

tion, black and light brown (weathered) colors, 1,5 km long and contains 12 layers (Fig. 6).

### Geochemical characteristics of oil shale and petroleum potential of deep-seated deposits

Eruption of mud volcanoes ends with emission of products at a depth of up to 6–8 km and more. These products are sole data, which contain information on geology of deep-seated sediments of mud volcanic areas, where have not been studied with drilling or detailed geophysical surveys. To study of hydrocarbon potential of these areas on the basis of geological and geochemical research of oil shale and oil-bearing rock association, which found in ejected products of mud volcanoes provides interesting results [3, 13–16].




Fig. 4. Schematic geological map and profile of Guba oil shale deposit




Fig. 5. Schematic map of oil shale containing areas, located between the rivers Velvelichay and Garachay




Fig. 6. Schematic geological map of Diyally oil shale deposit

Almost all mud volcanoes, located in Gobustan ejects oil shale and oil-bearing rock during their eruptions (Fig. 7).



Fig. 7. Oil shale (ejected by mud volcano in Gobustan)

There are more than 120 mud volcanoes in Shamakhi-Gobustan region (Fig. 8). Approximately all the mud volcanic structures in the region are potentially oil-bearing. Duvanny oil and gas field is developing in Southern Gobustan (Productive series, Miocene). Commercial oil and gas content of the productive series V and VII horizons has been established (over the Garadag break-down suite) at the Dashgil mudvolcanic area; Kyanizadag gas condensate field has been discovered in the Productive series; commercial oil influx from the Miocene sediments (Chockrak horizon) has been obtained on the Dashgil structure.

A geochemical study aimed at obtaining new information on stratigraphic intervals and depths of hydrocarbon in the study area to clarify the hydrocarbon generation and accumulation potential of the Eocene and the Miocene sediments, associated with mud volcanoes. The studies of oil shale, which sampled from mud volcanic products show that its organic composition ranges between 7,56–42,55 %.

According to data, obtained from a result of the extraction of oil shale, the largest number of soluble organic matter is observed in the composition of the mud volcano Gushchu – 8,34% (composition of kerogen – 34,02 %). Relatively fewer soluble organic matter is observed in the composition of oil shale, sampled from mud volcanoes Chapilmysh and Cheildag (0,56%). Most of the organic matter in the composition of oil shale samples accumulates in their kerogen portion (Table 1).

Formation and distribution of organic substances, mainly depend on the processes taking place at the final stages of diagenesis and the specific properties of the initial substances. In order to explain the process of natural catagenesis, held thermal laboratory analysis of oil shale. Thermolysis of samples, taken from various zones of mud volcanic areas, shows their Tmax. Samples of mud volcano Gushchu, rich with organic matter burn with several stages at temperatures of 300, 400 and 500 °C. Samples of mud volcanoes Chapilmish and Veys, containing relatively greater amounts of organic substances lose their masses at a temperature of 400 and 200–400 °C (Fig. 9, 10).






Fig. 8. Location map of mud volcanoes in Shamakhi-Gobustan region:

- 1 – Sarsura; 2 – Zeiva; 3 – Bizlan; 4 – Demirchi; 5 – Gyzmeidan; 6 – Yailag – Tudar;
- 7 – Gasymkend; 8 – Kekhnagyady; 9 – Kemchi; 10 – Kurkachidag; 11 – Hajly;
- 12 – Khilmilli; 13 – Garayaz; 14 – Agdere; 15 – Shikhandag; 16 – Nohur;
- 17 – Garanohur; 18 – Madrasa; 19 – Sarabil; 20 – Kyalakhana; 21 – Osmanbeili;
- 22 – Charhan; 23 – Nyuidi; 24 – Melikchobanly; 25 – Gyrlyg – Geoglyar;
- 26 – Chyrarly; 27 – Akhar – Bakhar; 28 – Jeirli; 29 – Chalov; 30 – Maraza;
- 31 – Gurbanchi; 32 – Nabur; 33 – Chaigur – banchy; 34 – Shimshadi;
- 35 – Kichik Maraza; 36 – Bozaakhtarma; 37 – Shikhzarli; 38 – Shorsulu; 39 – Ekakhana;
- 40 – Makhhlajik; 41 – Arabgadim; 42 – Juan; 43 – Gaiblar; 44 – Yeldarasi; 45 – Garajyuzlyu;
- 46 – West Tuva; 47 – East Tuva; 48 – South Tuva; 49 – Siyaki; 50 – west Veis;
- 51 – East Tuva; 52 – Neftik; 53 – Jengi; 54 – Syungur; 55 – Iyimish; 56 – Birgut;
- 57 – Donguzdug; 58 – Baygushlu; 59 – Sarydash – Bayanata; 60 – Gyrdag;
- 61 – Pirekeshkul MV group; 62 – Agdag; 63 – Arbat; 64 – Gyrgyshlag;
- 65 – Boransyz – Jylga; 66 – Agzygyr; 67 – Garyja; 68 – Charani; 69 – Chapilmish;
- 70 – Shakhgaya; 71 – Chukhuroglybozu; 72 – Gazanagyl; 73 – Sheitanud MV group;
- 74 – Gushchu; 75 – Kolany; 76 – Baidar; 77 – Ayazakhtarma; 78 – Ilkhychy;
- 79 – Sheikh Novruz; 80 – Sundi; 81 – Nardaranakhtarma MV group; 82 – Kyurdamich;
- 83 – Suleymanakhtarma; 84 – Cheilakhtarma; 85 – Gadridere; 86 – Hajiveli; 87 – Agnohur;
- 88 – West Cheildag; 89 – East Cheildag; 90 – Galandarakhtarma; 91 – Umbaki;
- 92 – West Davalidag; 93 – East Davalidag; 94 – Utagi; 95 – Agtapa; 96 – Beyuk Kyanizadag;
- 97 – Goturlug; 98 – Gylynych; 99 – Toragay; 100 – Kichik Kyanizadag; 101 – Hajiveli;
- 102 – Dashmardan; 103 – Shekikhan; 104 – Agdam MV group; 105 – Arzani; 106 – Durandag;
- 107 – Gotur; 108 – Agtirme; 109 – Emjek – emjek; 110 – Solakhay; 111 – Oyoug;
- 112 – Gyogyarchin; 113 – Dilyangyaz; 114 – Dashgil; 115 – Bala Bahar; 116 – Bahar;
- 117 – Garakyura; 118 – Airanteken; 119 – Saryboga; 120 – Goturdag

Table 1

Extraction of oil shale, sampled of mud volcanoes in Gobustan

Mud volcano	Amount of dissolved organic matter, %		Amount kerogen, %
	Chloroform	Alcohol-benzol (1:1)	
Shikhzarli	0,44	0,97	6,15
Pirekeshkul	1,00	0,61	19,97
Veys	0,51	0,42	16,85
Gushchu	2,54	5,80	34,02
Chapilmish	0,34	0,22	27,31
Suleymanakhtarma	1,12	0,62	8,75
Cheildag	0,18	0,38	9,35
Galandarakhatarma	1,04	0,74	6,12
Shekikhan	0,59	0,50	10,36
Agtirme	1,16	0,48	8,47
Solakhay	0,82	1,05	9,88
Dashgil	0,72	0,56	10,02
Ayrantoken	0,93	1,02	8,96
Durandag	1,01	0,52	12,35
Gotur	1,22	0,55	10,06




Fig. 9. Thermolysis of oil shale, sampled from mud volcanoes Shikhzarli, Suleymanakhtarma and Gushchu




Fig. 10. Thermolysis of oil shale, sampled from mud volcanoes Veys, Pirekeshkul and Chapilmish



**Table 2**

The two-stage pyrolysis of oil shale, sampled from mud volcanoes in Gobustan

Mud volcano	500–550°C			800–850°C		
	Water, %	Bitumen, %	Gas, %	Water, %	Bitumen, %	Gas, %
Shikhzarli	0,24	3,21	3,13	0,84	–	2,64
Pirekeshkul	0,42	4,12	4,53	0,32	0,42	2,81
Veys	4,25	5,60	6,31	–	–	1,19
Gushchu	1,35	6,18	17,69	0,32	–	11,77
Chapilmish	0,12	1,20	24,0	1,27	–	2,81
Suleymanakhtarma	0,81	3,87	2,34	0,43	–	2,27
Galandarakhatarma	0,25	4,15	2,12	0,33	–	2,11
Dashgil	1,80	3,12	7,01	–	–	0,80
Durandag	2,78	3,38	4,17	–	–	1,30
Gotur	0,56	0,45	10,67	–	–	5,53

Pyrolysis of oil shale samples of different mud volcanoes (Veys, Gushchu, Pirekeshkul, Galandarakhatarma and et al.) showed that up to 500°C temperature obtains fractions, similar to oil, but at higher temperatures, only gases. A sharp changing is observed at a temperature of 400°C (Table 2).

According to the amount of products obtained from oil shale, is making possible to define the structure of organic matter. Greater amount of bitumen indicates the structure of an aliphatic but others – like gas and light hydrocarbons an aromatic.

In addition to oil shale, oil-saturated rocks are also appear on the mudvolcanic crater zones after the eruptions of mud volcanoes [8]. In contrast to oil shale, they are mainly related to younger sediments (Miocene). The fraction, obtained by extraction (chloroform) mainly consists of asphaltenes (42,52–47,15%), and oil (26,86–37,96%).

Data obtained from chromatographic analysis (silica gel ASM) of oil fractions show that the majority of bitumen consist of monocyclic hydrocarbons (56,0–75,50%). The compounds similar to oil, but methane-naphthene relatively less than aromatic hydrocarbons (11,21–18,06%). The paraffins are almost absent in studied fractions.

### Conclusions

The analysis of oil shale objects in Azerbaijan shows that greater manifestations are developed in Gobustan and Absheron, during the Middle Eocene (Kone) and the Upper Miocene (Meotian). But Diyally oil shale deposit belongs to Sarmatian age. Regarding the conditions of formation of oil shale (within the shale facies) in these areas, it should be noted that a relatively significant manifestation (sustained by length and thickness) observed in

large synclines or trough: in the northern areas of Gobustan they trace in the cores of synclines (Charkishlak, Ambizlar, Shahandag, Agburun and etc.), in the southern areas, in the wings of synclinals (Boyuk Siyaki, Kichik Siyaki, Mayash, Jangidag, Islamdag and etc.) and in the front wings of anticlines (Buransiz-Bayanata, Jangichay and etc.).

The main practical features of Azerbaijan oil shale relate to its distribution regularities along the areas. Connected with different geological age, oil shale manifestations (Kichik Siyaki, Boyuk Siyaki, Islamdag, Baygushkaya, Uchtepe etc.) have found development within the same area (mainly in Gobustan) and closely spaced from each other. Such a regularities creates favourable conditions on joint development of these manifestation for their future exploitation. To locate away from the tracts of forest and the housing unit, is extra superiority of Azerbaijan oil shale from an economic and environmental point of view.

Analysis of thermolysis process shows that the maximum interval of mass loss for oil shale depends on the structure of organic matter, initial substances, geological conditions and thermocatalytic impacts and etc. Thus, if organic substances lose mass at high temperatures correspond an aliphatic, at low temperatures an aromatic type of structures.

The results of the pyrolysis show that the development of some kerogens of oil shale (found in ejected products of Chapilmish, Gotur and others mud volcanoes) has been weak in the oil generation processes, but others (Veys, Pirekeshkul, Galandarakhatarma and others) more intensive. Thus, if at low temperature the amount of obtained bitumen is much in former, but for the second group of mud volcanoes it almost absent. If compare the amount of gases, they are much greater (24,0%) in the samples of

Chapilmish mud volcano. The analytical study of such rocks show that related to re-changing of organic matter: long aliphatic C–C bonds are destroyed at the stage of ketogenesis, minerals and organic substances are separated from each other in kerogen composition and the whole process ends with formation of hydrocarbon.

IQ-spectrum of kerogen indicates its similarity to oil asphaltene and coal. In this regard, during the pyrolysis of kerogen (matured) obtains hydrocarbons. Thus, the results of laboratory tests show that some kerogens of oil shale (including of Chapylmysh mud volcano) are not yet fully matured.

Geological and geochemical analysis of oil shale, related to Paleogene-Miocene sediments and traces with a thickness of 3,5–4,5 km in the Central Gobustan (the location zone of Veys, Pirekeshkul mud volcanoes) provides the economic prospects in terms of the exploitation of shale gas.

The territory of South Gobustan is considered to be more promising in hydrocarbon generation, related to maturity of kerogen in oil shale (Eocene and Maikopian age). In addition, the comparative analysis of bitumen, which obtained from kerogen of oil shale (Eocene and Maikopian age) and oil-bearing rocks (Miocene age) confirms their close genetic relationship. The results confirm the possibility formation of hydrocarbons in the sediments of Eocene age (probably also accumulation in its granular and fractured reservoirs) and Maikopian series in the study area and their migration to the relatively younger Miocene reservoirs.

### References

1. Abbasov O.R. Geological and geochemical features of combustible slates of Gobustan (Azerbaijan) and their forecast inventories // Bulletin of the Atyrau Institute of oil and gas. – 2008. – № 2 (14). – P. 22–29.
2. Abbasov O.R., Mamedova A.N., Guseynov A.R., etc. Some new data of geochemical researches of combustible slates of Azerbaijan // Geology, geophysics and development of oil and gas fields. – 2013. – № 2. – P. 32–35.
3. Abbasov O.R., Ibadzade A.D., Hasayeva A.B., etc. Hydrocarbon potential of the deep-shipped deposits of Gobustan (Azerbaijan) (on the basis of combustible slates and oil-bearing breeds, emissions of mud volcanoes) // Resursovosproizvodyashchy, low-waste and nature protection technologies of development of a subsoil. – Kyrgyzstan, Bishkek, 2015. – P. 342–443.
4. Ali bakh A.A., Akhmedov G.A., Zeynalov M.M. Combustible slates of the Miocene of Azerbaijan // ANH. – 1962. – № 1. – P. 5–8.
5. Aliyev Hell. And., A.A's Bayrams. Some aspects of tectonics of mud-volcanic zones of Gobustan // Izv. AN of Azerbaijan, Science about Earth. – 2000. – № 1. – P. 129–131.
6. Aliyev Hell.And., Belov I.S., Bayrams T.A's. Combustible Paleogen-Miocene slates of Azerbaijan // Geologist of Azerbaijan. The scientific bulletin. – 2003. – № 8. – P. 68–80.
7. Aliyev Hell. And., Belov I.S., Aliyev G.A. Combustible slates of the Miocene of Azerbaijan // ANH. – 2000. – № 5. – P. 7–11.
8. Babayev F.R., Abbasov O.R., Mamedova A.N. etc. Studying of bitumens of Azerbaijan // Actual problems of humanitarian and natural sciences. – 2013 7 (54), 1 Is frequent. – P. 40–42.
9. Sultanov A.D., Sultanov R.G. Combustible slates of a southeast tip of Greater Caucasus Range in Azerbaijan // Az is THIS. Soviet Socialist Republic. – 1945. – T. 1. – № 1. – P. 11–13.
10. Sultanov A.D., Sultanov R.G. Materials on combustible slates of Azerbaijan // Izv. AN of Azerb. Soviet Socialist Republic. – 1947. – № 2. – P. 18–20.
11. Abbasov O.R. Oil Shale of Azerbaijan: Geology, Geochemistry and Probable Reserves // IJRSET – International Journal of Research Studies in Science, Engineering and Technology. – 2015. – Vo. 2, Issue 10. – P. 31–37.
12. Abbasov O.R. Distribution regularities of oil shale in Azerbaijan // ISJ Theoretical & Applied Science. – 2016. – Vol. 35, Issue 3. – P. 165–171.
13. Abbasov O.R., Akhundov R.V. Petroleum potential of Paleogene and Miocene deposits in Gobustan based on oil shale products of mud volcanoes // Baku World Forum of Young Scientists. – Baku, 2014. – P. 27–28.
14. Abbasov O.R., Baloglanov E.E., Akhundov R.V. Geochemical analysis of oil shale and oil-bearing rocks of Gobustan mud volcanoes // 6th International Conference of Young Scientists and Students "Multidisciplinary approach to solving problems of geology and geophysics" (Baku, October 12–15). – Baku, 2015. – P. 118–119.
15. Abbasov O.R., Baloglanov E.E., Akhundov R.V. Organic compounds in ejected rocks of mud volcanoes as geological and geochemical indicators: a study from Shamakhi-Gobustan region (Azerbaijan) // International Multidisciplinary Forum "Academic Science Week-2015". – Baku, 2015. – 02–04 November, P. 3–4.
16. Abbasov O.R., Ibadzadeh A.C., Mammadova A.N. Hydrocarbon generation potential of the deeper sediments of Azerbaijan // Integrated approach for unlocking hydrocarbon resources. – Baku, 2012. – P. 48.